

Theory of heating of hot magnetized plasma by Alfvén waves. Application for solar corona

T.M. Mishonov,^{*} M.V. Stoev,[†] and Y.G. Maneva[‡]

*Department of Theoretical Physics, Faculty of Physics,
University of Sofia St. Clement of Ohrid,
5 J. Bourchier Boulevard, BG-1164 Sofia, Bulgaria*

(Dated: February 5, 2008)

The heating of magnetized plasma by propagation of Alfvén waves is calculated as a function of the magnetic field spectral density. The results can be applied to evaluate the heating power of the solar corona at known data from satellites' magnetometers. This heating rate can be incorporated in global models for heating of the solar corona and creation of the solar wind. The final formula for the heating power is illustrated with a model spectral density of the magnetic field obtained by analysis of the Voyager 1 mission results. The influence of high frequency dissipative modes is also taken into account and it is concluded that for evaluation of the total coronal heating it is necessary to know the spectral density of the fluctuating component of the magnetic field up to the frequency of electron-proton collisions.

PACS numbers: 52.35.Bj, 52.50.Sw, 96.60.P-, 94.30.cq

I. INTRODUCTION

The discovery of the spectral lines of highly ionized iron in the spectrum of the solar corona showed in 1939 that the corona is much hotter than the radiating surface of the Sun. Why it is hotter by about two orders of magnitude is still a very discussed problem, which survived not only the Second World War, but also the whole development of the modern astrophysics and satellite technologies in the past half century. Definitely, this is the oldest still living mystery in front of the contemporary physics.

On the other hand, heating of the solar corona is a phenomenon in the simplest physical system - a hot ideal gas of electrons and protons. All the phenomena in a completely ionized hydrogen plasma are well-known and now it is a matter of a carefully proposed complementary set of experiments to arrive at the final conclusion. According to the second law of thermodynamics each process in the plasma surrounding the Sun produces some entropy and can be considered as a heating mechanism of the solar corona. At least one new idea per year appears under the Sun, concerning the heating of its corona. In a review article on the subject one can find more than a hundred references. Often they are being classified as *AC* and *DC* models¹ depending on whether a direct or an alternating electric current heats the plasma. But the coronal plasma can not be heated by an electric current, as it was proved thirty years ago⁸: "...it is difficult to understand how a cooler gas (the electrons) can heat a hotter gas (the protons) within a region so far away from the Sun, which is the ultimate source of most of the energy in interplanetary space", see². As far as the proton temperature is higher than the electron one $T_p > T_e$ the heating mechanism acts on the protons and the viscose friction of the shear flow with a velocity field $\mathbf{V}(z, t)$ is a natural explanation for the bulk power Q of the dissi-

pated energy⁵

$$Q = \frac{1}{2} \eta \sum_{i,j=1,2,3} (\delta_i V_j + \delta_j V_i)^2, \quad (1)$$

for $\text{div} \mathbf{V} = 0$. Such a heating process can be observed in every turbulent fluid. Imagine a waterfall for which the amplitude of turbulent pulsations is smaller than the velocity of sound and as a result the related heating after the calming of the turbulent motion is smaller $\Delta T/T \ll 1$. That is why the turbulence in the solar plasma cannot give an augmentation by two orders of magnitude of the coronal temperature. For an effective heating mechanism we need an effective mechanism for the energy transport.

This idea is supported by the recent development of the solar physics related to the random motion of magnetic footprints. The creation of Alfvén waves by the stochastic convection in the Sun is a very interesting hydrodynamic problem for the complete heating scenario (for additional references see Ref. 3). The purpose of the present work is to analyze what kind of local satellites' measurements should be done in order to confirm or reject the Alfvén wave mechanism for the coronal heating. Nowadays astrophysicists are able to use the available data provided by SOHO, TRACE, ACE, WIND in order to probe their model. However, the main defect of these experiments is the uncompleted set of data and that is why as far as the theory is concerned all the gladiators remain friendly sleeping in the arena. In the next section we will systematize the formulas for Alfvén waves from well-known textbooks and in the final section we will analyze what kind of research has to be done.

II. BASIC NOTIONS. MODEL EVALUATION

Calculation of the dissipated power density is an indispensable ingredient in every theory which claims to

reveal the coronal heating mechanism. The real scientific problem is to explain the heating power, not the local temperature jump. The local temperature is only a parameter to be used in a realistic global model for the solar corona, which includes the acceleration of the solar wind by an expansion of the heated plasma and all processes for the local energy balance.

We choose the z-axis oriented by the constant external magnetic field $\mathbf{B}_0 = \mathbf{e}_z$. We will investigate an Alfvén wave, propagating on this axis. We can choose the y-axis to be perpendicular to the external magnetic field and the wave vector $\mathbf{k} = k_x \mathbf{e}_x + k_z \mathbf{e}_z$. The Alfvén waves are linearly polarized and the wave component of the magnetic field $\mathbf{B} = \mathbf{B}_0 + \mathbf{B}'$ is oriented along the y-axis

$$\mathbf{B}'(\mathbf{r}, t) = B'_y \mathbf{e}_y, \quad B'_y = B_{y,0} \cos(\mathbf{k} \cdot \mathbf{r} - \omega_A t) e^{-\gamma z}, \quad (2)$$

where the null index denotes the wave amplitude, ω_A is the frequency of the Alfvén waves and the amplitude decrement γ is supposed to be small enough. For the wave period averaged density of the magnetic energy we have

$$w_B(z) = \frac{B_{y,0}^2}{2\mu_0} \frac{1}{2} e^{-2\gamma z}, \quad (3)$$

where $\mu_0 = 4\pi$ or $4\pi \cdot 10^{-7}$ in the practical system.

If we follow the evolution of a wave-packet, the density of the total energy (magnetic and kinetic) decreases exponentially over time

$$w(t) = \frac{B_{y,0}^2}{2\mu_0} e^{-t/\tau_A}, \quad (4)$$

where the absorption rate of the Alfvén waves

$$\frac{1}{\tau_A} = (\nu_{\text{kin}} + \nu_{\text{mag}}) k^2, \quad \nu_{\text{kin}} = \frac{\eta}{\rho}, \quad \nu_{\text{mag}} = c^2 \varepsilon_0 / \sigma \quad (5)$$

is given by a sum of the kinetic ν_{kin} and the magnetic ν_{mag} viscosities, expressed in terms of the plasma viscosity η , density ρ and the electrical conductivity of the fluid σ ; $\varepsilon_0 = 1/4\pi$ or $\varepsilon_0 = 1/\mu_0 c^2$ in the practical system.

For an order of magnitude evaluation of the amplitude decrement⁴ we will consider only the Alfvén waves $\omega_A = V_A k_z$ propagating along the external magnetic field lines $\mathbf{k} = k \mathbf{e}_z$ and discard the negligible influence of the resistivity ν_{mag}

$$\frac{1}{\tau_A} = 2\gamma V_A = \frac{\omega_A^2}{V_A^2} \frac{\eta}{\rho} = \frac{(2\pi f)^2}{V_A^2} \nu_{\text{kin}}, \quad (6)$$

where $f = \omega/2\pi$ is the circular frequency. The relation between the velocity of the Alfvén waves and the magnetic field can be found in any textbook on magnetohydrodynamics

$$\frac{1}{2} \rho V_A^2 = \frac{1}{2\mu_0} B_0^2, \quad \mathbf{V}_A = \frac{\mathbf{B}_0}{\sqrt{\mu_0 \rho}}, \quad \mathbf{B}_0 = B_0 \mathbf{e}_z. \quad (7)$$

We presume small fluctuations from the constant magnetic field so that the maximal distortion angle of the magnetic field lines

$$b_{y,0} = \frac{B_{y,0}}{B_0} \ll 1 \quad (8)$$

is also small and we can use the linear theory of Alfvén waves, propagating along the z-axis, whose phase velocity V_A coincides with their group one v_{gr}

$$\omega_A(\mathbf{k}) = |\mathbf{V}_A \cdot \mathbf{k}|, \quad \omega_A = V_A k_z, \quad k_z > 0, \quad v_{gr} = V_A. \quad (9)$$

For the bulk power density of the dissipated wave energy we obtain

$$Q = \frac{w}{\tau_A} = \frac{B_{y,0}^2}{2\mu_0 \tau_A}. \quad (10)$$

In this model evaluation we have formally taken $\mathbf{r} = 0$, assuming that the weak space dependence of the magnetic field is included in the local value of the wave amplitude,

$$B'_y(t) = B_{y,0}^{(\omega)} \cos(\omega t), \quad (11)$$

measured by an AC satellite's magnetometer. Here the upper index (ω) emphasizes that up to now we have considered a monochromatic wave whereas for the calculation of the total absorption rate we have to perform a frequency summation over the whole Alfvén waves spectrum. In reality we ought to take into account also the influence of the Doppler effect, caused by the solar wind, but as a gedanken-experiment let us consider, that the frequency is measured in a system, bound to the coronal plasma. Imagine that the magnetometer has a frequency-filter and detects only frequencies in a narrow range (f_1, f_2) , where $f_2 - f_1 \ll f_1$. The most important characteristic of the Alfvén waves is their spectral density¹³, given by a sum of the wave energies in a narrow frequency interval

$$\Phi(f) = \frac{1}{f_2 - f_1} \sum_{f \in (f_2, f_1)} \left(B_{y,0}^{(f)} \right)^2, \quad f = \frac{\omega}{2\pi}. \quad (12)$$

We can assume that this spectral density is given by an analogue filter, but in fact we have to do a Fast Fourier Transformation (FFT) for the magnetic field as a function of time. Generally speaking $\Phi(f)$ is a tensor, but for an illustrative purpose let us investigate only the y-components of the Alfvén waves. Then the heating power is given by a frequency integral of the damping rate and the spectral density taken up to the maximal frequency for which the magnetohydrodynamic approach is still applicable, i.e. the maximal electron-proton collisions frequency $\omega_{\text{max}} = 1/\tau_e$:

$$Q_{\text{abs}} = \frac{1}{2\mu_0} \int_0^{1/2\pi\tau_e} \frac{\Phi(f)}{\tau_A(f)} df. \quad (13)$$

In the damping rate we have to substitute the kinematic viscosity with that of a hydrogen plasma⁶

$$\nu_{\text{kin}} = \frac{\eta}{\rho} = 0.4 \frac{T_p^{5/2}}{M^{1/2} e^4 n_p L_p}, \quad (14)$$

where

$$\frac{1}{a^2} = 4\pi e^2 \left(\frac{n_e}{T_e} + \frac{n_p}{T_p} \right), \quad e^2 \equiv \frac{q_e^2}{4\pi\epsilon_0} \quad (15)$$

is the Debye radius, T_p and T_e are respectively the proton and electron temperatures multiplied by the Boltzmann constant, and

$$L_p = \ln \frac{T_p a}{e^2} \gg 1 \quad (16)$$

is the Coulomb logarithm as the thermal velocity of the protons is much smaller than the Bohr's velocity

$$v_{T_p} \ll v_{\text{Bohr}} \equiv \frac{e^2}{\hbar}, \quad M v_{T_p}^2 = T_p. \quad (17)$$

Here we have used the elementary gas-kinetic evaluation for the viscosity

$$\eta \sim n_p M l_p v_{T_p}, \quad (18)$$

as well as the mass density of the plasma, which is practically determined by the density of the protons n_p and their mass M

$$\rho = M n_p. \quad (19)$$

In the derivation of the upper evaluation we have supposed that two protons with kinetic energy T_p can be brought to a minimal distance r_{min}

$$\frac{e^2}{r_{\text{min}}} \sim T_p. \quad (20)$$

This distance parameterizes the effective cross-section ς_p for the proton-proton collisions, which participates in the evaluation for the mean free path of the protons

$$l_p \varsigma_p n_p = 1. \quad (21)$$

The Coulomb logarithm additionally increases the cross-section ς_p

$$\varsigma_p \sim \pi (2r_{\text{min}})^2 L_p. \quad (22)$$

For the hot coronal plasma the kinematic viscosity is much bigger than the magnetic one $\nu_{\text{kin}} \gg \nu_{\text{mag}}$. This determines the diffusion of the magnetic field due to the ohmic resistance.

For evaluating the maximal frequency at which magnetohydrodynamics can still be applied we have to calculate the maximal frequency at which the density of the electric current, transferred by the electrons, $\mathbf{j}$ creates

through the magnetic field $\mathbf{B}$ bulk Lorentz force, acting on the proton fluid with mass density ρ . This is in fact the frequency of the electron-proton collisions $1/\tau_e$, which we take from the elementary kinetic gas theory. We can use the duration of the mean free path of electrons τ_e to determine the electric conductivity of the plasma

$$\frac{\sigma}{4\pi\epsilon_0} \sim \frac{e^2 n_e}{m} \tau_e. \quad (23)$$

The mean time between two electron collisions is defined by the electron mean free path l_e and their thermal velocity v_{T_e}

$$\tau_e = \frac{l_e}{v_{T_e}}. \quad (24)$$

The relation between the electron density n_e , their free path l_e and effective cross-section ς_e is the same as for protons

$$l_e n_e \varsigma_e = 1, \quad \varsigma_e \sim \pi (2r_{\text{min}})^2 L_e, \quad (25)$$

only the expression for the Coulomb logarithm should be changed

$$L_e = \ln \frac{T_e}{\hbar \omega_{pl}} \gg 1, \quad (26)$$

as even the velocity of the non-relativistic electrons is much greater than the Bohr velocity

$$\frac{e^2}{\hbar} \ll v_{T_e} \ll c. \quad (27)$$

The plasma frequency ω_{pl} is much higher than the collisions frequency

$$\omega_{pl}^2 = \frac{n_e e^2}{m}, \quad (28)$$

and yet it is located in the radio-range.

These formulas give the well-known evaluations for the electron-proton collisions frequency

$$\frac{1}{\tau_e} = \frac{4\pi e^4 n_e L_e}{m^{1/2} T_e^{3/2}} \sim \frac{v_{T_e}}{L_e} \sim \frac{1}{\nu_e} \sim \omega_{A,c} \sim 2\pi f_c, \quad (29)$$

as well as for the conductivity of the fully ionized hydrogen plasma

$$\frac{\sigma}{4\pi\epsilon_0} = 0.6 \frac{T_e^{3/2}}{e^2 m^{1/2}}. \quad (30)$$

These model kinetic evaluations applied to the coronal plasma could be used for calculation of the heating power due to absorption of Alfvén waves. For an analytical illustration we will utilize the interpolation formula, used for analysis⁷ of the magnetometric data from Voyager 1

$$\Phi(f) \sim \frac{\mathcal{D}}{f^2}. \quad (31)$$

Such a spectral density $1/f^2$ is naturally explained in the framework of a stochastic Langevin MHD, as Mishonov and Maneva pointed out³, Eq. 41, and it is not related to any phase-coherent structures which could be distinguished from incoherent turbulent fluctuations, cf. with the work by Roberts and Goldstein¹¹. The Kolmogorov turbulence $f^{-5/3}$ gives^{9,10} a slightly (17%) different power. The idea that convective granules produce a continuous stream of noise generating waves is actually very old¹⁴. We believe that this is a common property of all turbulent magnetized plasmas and we suggest the hypothesis that the $1/f^2$ spectral density could be used even for the physics of accretion disks.

Bearing in mind all above, the general formula Eq. (13) gives the final model evaluation for the bulk heating power

$$Q_{\text{abs}} = 0.4(2\pi)^2 \frac{\mathcal{D}}{B_0^2} \left(\frac{M}{m}\right)^{1/2} \frac{L_e}{L_p} \left(\frac{T_p}{T_e}\right)^{3/2} n_p T_p. \quad (32)$$

This heating rate actually sets the upper limit for the absorbed bulk power that can be derived in the magnetohydrodynamic framework when we take into account the influence of the diffusion Alfvén modes with frequencies $\omega > 1/\tau_A$ as well. However, these diffusion modes are able to heat only the narrow “skin layer” above the limb. Far away from the Sun (at the height of the satellites) we have to account only for those Alfvén waves, propagating with $\omega_A > 1/\tau_A$. If we substitute the proton-electron collisions frequency cut-off $1/\tau_e$ in Eq. (13) with the Alfvén waves damping rate we will end up with a different assessment for the heating rate

$$\tilde{Q}_{\text{abs}} = \frac{\pi \mathcal{D}}{\mu_0}. \quad (33)$$

Hereby we obtain another time constant related to kinetics of the protons that is rather significant for the theory of coronal plasma heating

$$\frac{1}{\tau_h} = \frac{\tilde{Q}_{\text{abs}}}{n_p T_p} = \frac{\pi \mathcal{D}}{\mu_0 n_p T_p}. \quad (34)$$

This time constant τ_h together with the velocity of the solar wind v_{wind} form a dimensionless parameter

$$\Upsilon = \frac{r_{\text{Sun}}}{v_{\text{wind}} \tau_h}, \quad (35)$$

which shows at what distance from the Sun r_{Sun} the discussed mechanism for coronal heating stays important and should be incorporated in the coronal dynamics; at $\Upsilon \ll 1$ the influence of the Alfvén waves heating is negligible. According to our statistical approach, the heating time-constant depends on the choice of the frequency cut-off. Therefore the role of the frequency cut-off for the coronal heating due to viscous damping of magnetohydrodynamic waves is extremely important and we wish to find out what kind of physical processes in the plasma prevail to determine it. We have to pay attention that

the ratio of these two cut-offs significantly varies with the solar radius, starting from a large number in the photosphere, where the collisions frequency dominates the Alfvén waves attenuation, and reaching the order of unity in the lower corona, where the plasma β becomes rather small

$$\frac{\tau_{A,c}}{\tau_e} = 0.4\pi\beta \frac{n_e L_e}{n_p L_p} \left(\frac{M T_p^3}{m T_e^3}\right)^{1/2}, \quad \beta = \frac{n_e T_e + n_p T_p}{(B_0^2/2\mu_0)}. \quad (36)$$

As one can see, the most important characteristic in the result for the absorbed power Eq. (32) is the spectral density of the angle of deviation of the magnetic field lines from their state of static balance position.

Due to this distortion as strings of a harp

$$(b^2)_\omega = \frac{\Phi(f)}{B_0^2} = \frac{(B_{y,0}^2)_f}{B_0^2} \approx \frac{\mathcal{D}}{B_0^2 f^2}, \quad (37)$$

we can listen to the impressive symphony of the convective turbulence of the Sun and its corona is heated mainly by the high-frequency part of the spectral density.

III. DISCUSSION

A new theory often means data mining and experimental data processing, but our purpose is not to extract the electron and proton temperatures from SOHO data; our purpose is to explain a local property of the magnetized plasma, pierced by Alfvén waves.

We have calculated the absorption power of the Alfvén waves Q_{abs} , that is the power transmitted per unit volume from the Alfvén waves to the protons. In order to incorporate this leading (according to our opinion) power in a coherent picture we ought to include Q_{abs} in the total plasma energy balance which takes into account all possible energy losses or gains: the work due to the gas expansion, the change of the temperature which is proportional to the density of the kinetic energy, the energy transmission from protons to electrons by collisions, bremsstrahlung of electrons, etc. In other words, Q_{abs} should be the last missing ingredient of an almost completed model for the coronal heating and creation of the solar wind.

Few words should be said about how possibly our scenario could be rejected; the main defect of the contemporary coronal investigations is that none of the proposed mechanisms is consigned to the dustbin. Even the magnetic reconnections which predict significant Ohmic heating and $T_e \gg T_p$. As we propose Alfvén waves as a heating mechanism the perpendicular to the static magnetic field pulsations of the velocity and the magnetic field should be in relation corresponding to Alfvén waves⁴ Eq. (69.11)

$$\frac{v_\alpha}{V_A} = -b_\alpha = -\frac{B'_\alpha(z, t)}{B_0}, \quad \alpha = 1, 2, 3. \quad (38)$$

Such a relation was observed during the first space missions¹². In other words the present heating scenario should be abandoned if the correlation coefficient between the velocity and the magnetic field is not high enough. According to our picture the waves are generated by the turbulence in the dense photospheric layer of the Sun where the influence of the magnetic field is relatively weak and mainly transversal incompressible magnetohydrodynamic waves (MHD) waves are generated. These waves reach the regions with negligible pressure (i.e. low plasma β regions) but still their polarization corresponds to MHD waves in an incompressible fluid.

IV. CONCLUSIONS

The model problem for Alfvén waves transmission from a magnetized turbulent half-space to a non-turbulent half-space has been recently solved³. This model approach can be further developed by an inclusion of the inhomogeneity of the magnetic field crossing the two half-spaces interface. However, for the coronal heating theory we consider as more appropriate to use experimental data for the Alfvén waves spectral density. But the accuracy of the magnetic field measurements is much higher than that of the tachometry of the solar wind - that is why we focus on magnetometric data for the spectral density of the magnetic field $\Phi(\omega)$. This tensor ought to have two equal eigen values and another smaller one related to the static field component. Of course, it would be much better to have data for the magnetic field and velocity correlation. Investigations of the general 6×6 matrix, including all correlation coefficients, should be planned for the forthcoming space missions. The most important problem is where the spectral density of the MHD waves comes to an end. From a microscopic point of view the cut-off frequency is at electron-proton collisions rate $1/\tau_e$ which is well in the range of the standard radio frequency measurements. However, the maximal frequency $\omega_{\max}$ should be determined by the concrete physical conditions. For example, the inverse minimal size of the solar granulation could also determine the cut-off wave-vector k_c . Another possibility for the maximal frequency is to be set by the waves propagation condition $\omega_A \tau_A > 1$, which results in

$$\omega_c = V_A k_c = \nu k_c^2 = V_A^2 / \nu = B_0^2 / \mu_0 \nu \rho, \quad k_c = V_A / \nu. \quad (39)$$

Above this frequency the waves are pure dissipative modes $\omega \approx i\nu k^2$ with frequency-dependent depth of the "skin layer" $\delta = \sqrt{\nu/\omega}$.

One of the problems we encounter today is that the high-frequency MHD waves are absorbed during the coronal heating and all the satellites' reports have a distorted spectral density like the red light of a sunset. In order to observe the bare MHD waves we need to have magnetometric data from a satellite-kamikaze, but this would be a too expensive toy to be planned and realized. We

need a more ingenious solution. We ought to analyze online which region of the Sun emits the MHD waves, investigated by the magnetometer.

As a real break-through of the solar physics we consider the investigation of the correlation between the stochasticity of the space magnetic field delivered by the MHD waves and the stochasticity of the solar surface, obtained by optical data. Correlation of SOHO Dopplergrams with magnetograms is an important task, which should be put on the agenda in the global magnetohydrodynamic modeling of the solar corona. Before this is done we can just extrapolate the spectral density from the experimentally accessible region to the cut-off frequency $1/\tau_e$, as it was done in the derivation of our model evaluation Eq. (13). This Q_{abs} can be compared with the necessary heating power of the global MHD models for the solar corona. This final formula contains only experimentally known parameters and can be used in the global MHD models for the solar corona.

We would like to close with a qualitative discussion of the proposed mechanism in this work. As it is well-known, the leap of two orders of magnitude of the coronal temperature is rather sharp, compared to the height to the optically dense layers. This leap is a kind of a domain border, typical for the condensed matter physics. This sharp border ought to have at least qualitatively a good explanation in each mechanism for heating of the coronal plasma and this should be used as an important aesthetic criterion for the applicability of each coronal heating model. For most of them the closer we are to the dense layers the more intense is the heating and the observed temperature jump is practically unexplainable. Let us discuss this leap in the framework of the proposed scenario: slightly above the optically dense layers, emitting black body radiation $\propto T^4$, the dense plasma is transparent for the MHD waves as the viscosity $\rho\nu \propto \eta \propto T^{5/2}$ is almost density independent. In this optically transparent layer the heating is negligible and the cooling is realized perhaps by a small residual convective turbulence. Imagine that all the plasma above is cool with $T \sim 6$ kK and let us consider its temperature stability. Imagine a local fluctuation - a small augmentation of the temperature. This increasing will lead to the enhancement of the viscosity η and the heat absorption from the MHD waves, piercing with an almost constant energy density through every layer of the plasma. This increasing of the heating leads to an extra-augmentation of the temperature, which cannot be compensated from the thermal conductivity in the rarefied plasma. Thus we have a positive feed back and the plasma explodes from the absorption of high-frequency MHD waves. This is simultaneously the mechanism explaining the creation of the solar wind. As a comparison let us mention that for none of the chemical explosions at room temperature we do not have an increasing of the temperature with as much as two orders of magnitude. It would be of a significant interest to investigate the leap of the spectral density of the magnetic field $\Phi(\omega)$ at this frontier.

As a conclusion we would like to mention that in fact the first historical explanation of the solar coronal heating problem is still on the arena, retrieving another life or at least an *impetus* from the current satellites' data.

Acknowledgments

Discussion and support by D. Damianov, R. Erdelyi, I. Rousev, T. Zaqarashvili and I. Zhelyazkov are highly

appreciated.

-
- * E-mail: mishonov@phys.uni-sofia.bg
 - † E-mail: martin.stoev@gmail.com
 - ‡ E-mail: yanamaneva@gmail.com
 - ¹ C. H. Mandrini, P. Démoulin, J. A. Klimchuk, *The Astrophysical Journal*, **530** 999-1015 (2000).
 - ² R. Schwenn, E. Marsch, *Physics of the Inner Heliosphere*, Vol. 2, p. 99 (1991).
 - ³ T. Mishonov, Y. Maneva, "Burgulence and Alfvén Waves Heating Mechanism of Solar Corona", astro-ph/0609609.
 - ⁴ L. Landau and E. Lifschitz, *Course of Theoretical Physics*, Vol. 8 *Electrodynamics of Continuous Media*, Sec. 69, problem.
 - ⁵ L. Landau and E. Lifschitz, *Course of Theoretical Physics*, Vol. 6 *Fluid Mechanics*, Eq. (16.3).
 - ⁶ L. Landau and E. Lifschitz, *Course of Theoretical Physics*, Vol. 10 *Physical Kinetics*, Sec 43.
 - ⁷ L. F. Burlaga, W. H. Mish, *J. Geophys. Res.*, **90**, 4373 (1985); **92**, 1261 (1987).
 - ⁸ Feldman, W.C., J.R. Asbridge, S.J. Bame, J. T. Gosling, "High-speed solar wind parameters at 1 AU", *J. Geophys. Res.* **81**, 5054-5060 (1976); R. Schwenn, "Large-Scale Structure of the Interplanetary Medium" in *Physics and Chemistry in Space - Space and Solar Physics*, Vol. 20 *Physics of the Inner Heliosphere* Vol. 1, Editors: R. Schwenn, E. Marsch, Springer-Verlag Berlin (1990).
 - ⁹ Burlaga, L. F., "Hydromagnetic waves and discontinuities in the solar wind" *Space. Sci. Review*, **12**, 600, (1971); E. Marsch, "MHD Turbulence in the Solar Wind" in *Physics and Chemistry in Space - Space and Solar Physics*, Vol. 21, *Physics of the Inner Heliosphere* Vol. 2, Editors: R. Schwenn, E. Marsch, Springer-Verlag Berlin (1991).
 - ¹⁰ Kolmogorov, A. N., "The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers, *Compt. Rend. Acad. Sci. U.R.S.S.*, **30**, 301, (1941); E. Marsch, "MHD Turbulence in the Solar Wind" in *Physics and Chemistry in Space - Space and Solar Physics*, Vol. 21 *Physics of the Inner Heliosphere* Vol. 2, Editors: R. Schwenn, E. Marsch, Springer-Verlag Berlin (1991).
 - ¹¹ Roberts, D. A., M. L. Goldstein, "Spectral signatures of jumps and turbulence in interplanetary speed and magnetic field data", *J. Geophys. Res.*, **92**, 10105, (1987); E. Marsch, "MHD Turbulence in the Solar Wind" in *Physics and Chemistry in Space - Space and Solar Physics*, Vol. 21 *Physics of the Inner Heliosphere* Vol. 2, Editors: R. Schwenn, E. Marsch, Springer-Verlag Berlin (1991), p. 168.
 - ¹² R. Bruno, B. Bavassano, U. Villante, "Evidence for long-period Alfvén waves in the inner solar system" *J. Geophys. Res.*, **90**, 4373, (1985); cf. with the theoretical reviews: A. Barnes, "Interplanetary Alfvénic fluctuations: A stochastic model" *J. Geophys. Res.*, **86** 7498 (1981); E. Marsch, "MHD Turbulence in the Solar Wind" in *Physics and Chemistry in Space - Space and Solar Physics*, Vol. 21 *Physics of the Inner Heliosphere* Vol. 2, Editors: R. Schwenn, E. Marsch, Springer-Verlag Berlin (1991).
 - ¹³ Bavassano, B., M. Dobrowolny, F. Mariani, N. F. Ness, "Radial evolution of power spectra of interplanetary Alfvénic turbulence" *J. Geophys. Res.*, **87**, 3617, (1982); E. Marsch, "MHD Turbulence in the Solar Wind" in *Physics and Chemistry in Space - Space and Solar Physics*, Vol. 21, *Physics of the Inner Heliosphere* Vol. 2, Editors: R. Schwenn, E. Marsch, Springer-Verlag Berlin (1991), Figs. 10.2-4, 10.19.
 - ¹⁴ E. Schatzman, *Annales d'Astrophysique*, **12**, 203 (1949); M. Schwarzschild, *Astrophysical Journal*, **107**, 1 (1948).